

Two New Combinations in Peruvian *Taltomata* (Solanaceae)

Author(s): Thomas Mione and Felix G. Coe

Source: Novon, Vol. 2, No. 4 (Winter, 1992), pp. 383-384

Published by: Missouri Botanical Garden Press

Two New Combinations in Peruvian *Jaltomata* (Solanaceae)

Thomas Mione

Biological Sciences, Copernicus Hall, Central Connecticut State University, New Britain,
Connecticut 06050-4010, U.S.A.

Felix G. Coe

Ecology and Evolutionary Biology, University of Connecticut, Storrs, Connecticut
06269-3043, U.S.A.

ABSTRACT. The new combination *Jaltomata weberbaueri* (Dammer) Mione is based on *Saracha weberbaueri* Dammer; this species resides in Peru, Dept. Ancash, between 3,300 and 3,700 m elevation. The new combination *Jaltomata aspera* (Ruiz & Pavón) Mione is based on *Atropa aspera* Ruiz & Pavón; *Hebecladus asperus* (Ruiz & Pavón) Miers is a synonym. This species resides in Peru, Dept. Lima, from near sea-level to 1,800 m, and is rare, likely due to habitat destruction by humans.

Jaltomata (tribe Solaneae) is a genus of about 28 species of tropical and subtropical perennial herbs and small shrubs that are distributed from Arizona, U.S.A., to southern Bolivia, and with one species each in the Greater Antilles and on the Galápagos Islands.

During study of the genus *Jaltomata* we have found the following new combinations to be necessary. The species treated here are similar but are easily distinguished. *Jaltomata weberbaueri* resides above 3,000 m and is glabrate, while *J. aspera* resides below 1,800 m and is pubescent and ciliate.

Jaltomata weberbaueri (Dammer) Mione, comb. nov. Basionym: *Saracha weberbaueri* Dammer, Bot. Jahrb. Syst. 37: 638. 1906. TYPE: Peru. Ancash: Prov. Cajatambo, Ocros, *A. Weberbauer* 2657 (holotype, B destroyed, photo of B sheet F neg. 2558, WIS).

Saracha weberbaueri Dammer subsp. *pallascana* Bitter, Repert. Spec. Nov. Regni Veg. 19: 267. 1924. *Saracha pallascana* (Bitter) Macbride, Publ. Field Columbian Mus., Bot. Ser. 8: 110. 1930. TYPE: Peru. Ancash: Prov. Pallasca, valley of the river Pelagatos, near Yungabal, *A. Weberbauer* 7225 (holotype, B destroyed, photo of B sheet F neg. 2559, WIS; isotype, F, photo of F sheet F neg. 49345, WIS).

See Macbride (1962, as *Saracha weberbaueri*) for an English description of this taxon.

Additional specimens examined. PERU. ANCASH: Prov. Recuay, km 107 between Recuay and Pativilca, (fl), *M. Dillon et al.* 3174 (BH, GH, MO); Prov. Aija, cuchu cuchu, entre Huamanwilca y Sequipoquian, *C. Ochoa & A. Salas* 15160 (NY).

Habitat. Grasslands with scattered bushes, *A. Weberbauer* 2657, 7225; moist ravines, *M. Dillon et al.* 3174.

Altitude. 3,300–3,700 m.

Phenology. Flowering and fruiting January through March.

Local name. “Tomentito,” *M. Dillon et al.* 3174.

Jaltomata aspera (Ruiz & Pavón) Mione, comb. nov. Basionym: *Atropa aspera* Ruiz & Pavón, Fl. Peruv., et Chilensis 2: 45. 1799. *Ulticoma aspera* (Ruiz & Pavón) Raf., Sylva Telluriana 55. 1838. *Hebecladus asperus* (Ruiz & Pavón) Miers, London J. Bot. 4: 322. 1845. TYPE: Peru. In collibus Limae moninatis Amancae, *Ruiz López & Pavón* s.n. (holotype, presumably MA not seen).

Saracha ciliata Miers, Ann. Mag. Nat. Hist., ser. 2, 3: 446. 1849, syn. nov. TYPE: Peru. Valley of Lima, Mathews 834 (holotype, K).

Saracha lacrima-virginis Bitter, Repert. Spec. Nov. Regni Veg. 17: 341. 1921, syn. nov. TYPE: Peru. Amancae, *A. Weberbauer* 1587 (holotype, B destroyed, photo of B specimen F neg. 2551, WIS).

Saracha urbaniana Bitter & Dammer, Repert. Spec. Nov. Regni Veg. 17: 342. 1921, syn. nov. TYPE: Peru. Above San Bartolomé, Lima-Oroya Railroad, *A. Weberbauer* 5262 (5262 = 5305 according to label on photo at WIS) (holotype, B destroyed, photo of B specimen F neg. 2557, GH, WIS; isotype, GH).

See Macbride (1962, as *Hebecladus asperus*, *Saracha ciliata*, and *S. urbaniana*) for English description of this taxon.

Additional specimens examined. PERU. LIMA: Prov. Huarochiri, valley of Rio Rimac, near Lima-Oroya highway at km 70 E of Lima, *T. H. Goodspeed & R. D.*

Metcalf 30248 (MO, US); near Lima, *W. Nation s.n.* (K, on same sheet as holotype of *S. ciliata*); hills of Mongomarca, *A. Weberbauer* 5235 (US). LIMA: Amancaes, *A. Weberbauer* 5246a (F, US), *A. Weberbauer* 5305 (F); San Geronimo, *J. F. Macbride* 5912 (F), *D. Stafford* 1932 (K).

Habitat. Loma-formation, *A. Weberbauer* 1587, 5235, 5246a; rocky hillside, *Macbride* 5912; dry open hillside, granite soil, with cacti, *Goodspeed & Metcalf* 30248; sandy hills of Lima, shady places, between stones, rare, *W. Nation s.n.*

All specimens examined were collected prior to 1943; perhaps endangered, due to habitat destruction by humans.

Altitude. Near sea-level to 1,800 m.

Phenology. Flowering and fruiting July through October, and March.

Local name. "Lacrima de la virgen," *A. Weberbauer* 1587.

Acknowledgments. We thank Gregory J. Anderson, Tilton Davis IV, William D'Arcy, Michael Nee, and the herbarium staffs at BH, BM, C, COLO, F, GH, K, LD, MERF, MO, NY, US, and WIS for loan of specimens and/or accommodating visits, Thomas Stark for assistance with German, and Armando T. Hunziker and Luis Bernardello, who suggested this group for study. Support was from fellowships from the Graduate School and the Research Foundation of the University of Connecticut to T. Mione, and an N.S.F. grant to T. Mione and Gregory J. Anderson.

Literature Cited

- Macbride, J. F. 1962. Solanaceae. Field Mus. Nat. Hist., Bot. Ser. 13, part V-B, No. 1. [*Jaltomata* as *Saracha* and *Hebecladus*.]